



- **FACT SHEET No. 9**

Joint Pain in Pet Dogs and Cats

[Duncan Lascelles, BVSc, PhD](#)

Joint pain, particularly associated with osteoarthritis (OA), is common in companion animals such as dogs, cats, and horses. It results in impairment in mobility and performance of activities and is associated with spontaneous and induced pain. In pet dogs, osteoarthritis is a common condition possibly affecting a quarter of the population. OA in dogs is considered to be very similar to human OA [1] and therefore a potential spontaneous disease model. [15] [7] Additionally, such a “spontaneous model” has the added benefit that these companion dogs share the same environment as humans, potentially making the model more relevant than some rodent models. [15]

In pet cats, radiographic evidence of OA/degenerative joint disease (DJD) is apparent in up to 90 percent of all cats, [14] with an estimated 50 percent of these having clinical signs of impairment due to joint pain. Less is known about the etiology of OA in cats than in dogs, but the degenerative disease process appears very similar to that of other species. [1, 9] Other painful degenerative joint diseases such as the immune-mediated arthropathies occur in dogs and cats and may be underdiagnosed.

Etiology and Pathophysiology

In contrast to humans, OA in dogs is mainly due to developmental orthopedic diseases—hip dysplasia, elbow dysplasia, osteochondrosis dissecans, non-traumatic cranial cruciate ligament degeneration—and so is considered an early onset disease and a lifelong disease. The joints most commonly affected are the hip, stifle, and elbow.

In cats, the etiology of OA or DJD is less well understood, but the degenerative process appears to be very similar to other species. The joints most commonly affected are the hip, stifle, tarsus, and elbow.



© Copyright 2016 International Association for the Study of Pain. All rights reserved.

IASP brings together scientists, clinicians, health-care providers, and policymakers to stimulate and support the study of pain and translate that knowledge into improved pain relief worldwide.

In both cats and dogs, all tissues of the joint are involved in the degenerative process, and pain is commonly associated with the disease. Although pain cannot be predicted by radiographic appearance, this procedure does predict alterations in range of motion. Peripheral and central nervous system plasticity associated with joint pain has been demonstrated in both cats [10] and dogs [18] and is thought to contribute to the overall pain state.

Pain associated with joint disease results in impaired or altered mobility, impaired ability to perform activities, and altered behavior. In dogs, it has been shown to disrupt sleep [11] and is thought to impair cognitive function. In both species, the multidimensional effects of pain appear similar to the effects in humans.

Clinical Signs and Diagnosis

In clinical veterinary practice, diagnosis is centered on four elements:

1. Owner-reported activity impairment. This is more readily detected in dogs, and several clinical metrology instruments have been developed to measure this (CBPI [4]; LOAD [17]) One owner-completed clinical metrology instrument has been developed to detect DJD-associated pain and activity impairment in cats (FMPI [2]).
2. Pain on manipulation of affected joints during an orthopedic evaluation, with pain being measured as a behavioral response.
3. Radiographic evidence of osteoarthritis (effusion; osteophytes; subchondral sclerosis; joint-associated mineralization).
4. Synovial fluid analysis.

Overall, outcome measures are relatively well developed in the dog but less so in the cat, which has influenced the development of therapies. In referral centers or in the comparative research setting, the effects of pain can be measured in both species by measuring limb use (kinetic variables measured using force plates or pressure sensitive walkways) [12], spontaneous activity through accelerometry [5], and central plasticity through quantitative sensory threshold testing. [6, 18]

Treatment

Because of a relative paucity of evidence-based information regarding dogs and cats, much of the current clinical approach to treatment is based on information borrowed from human medicine.

Dogs:

- A multimodal drug and non-drug approach is recommended to manage OA-associated pain. [8]



© Copyright 2016 International Association for the Study of Pain. All rights reserved.

IASP brings together scientists, clinicians, health-care providers, and policymakers to stimulate and support the study of pain and translate that knowledge into improved pain relief worldwide.

- The only class of drug approved by the U.S. Food and Drug Administration Center for Veterinary Medicine is the NSAID class (several approved), though other classes (caninized anti-nerve growth factor and prostaglandin E4 receptor antagonists (piprants), among others, are in development).
- Local (intra-articular) therapies are occasionally used, and several drugs (such as capsaicin analogues) are in development.
- Most evidence exists for efficacy of NSAIDs, dietary modulation through omega-3 fatty- acid supplementation, [16] weight management, and exercise.
- Adjunctive drug therapy (amantadine, tramadol, gabapentin) is commonly employed (evidence for efficacy of amantadine; [13] limited evidence for oral tramadol, which is metabolized very differently in the dog; no evidence for gabapentin). Overall, there is a lack of studies evaluating these therapies.
- Physical rehabilitation (exercise and other physical modalities) is commonly employed.
- Surgical joint replacement is available and used in the dog (hip, stifle, and elbow).
- Combination steroid and immunosuppressive therapies are used in the management of immune-mediated joint pain.

Cats:

- A multimodal drug and non-drug approach is recommended to manage OA and DJD-associated pain. [3]
- There are no FDA-approved drug treatments and only one approved drug in the European Union (NSAID), though other classes (felinized anti-nerve growth factor and prostaglandin E4 receptor antagonists (piprant), among others, are in development).
- Most evidence exists for efficacy of NSAIDs and dietary modulation through omega-3 fatty-acid supplementation. [16]
- Adjunctive drug therapy (mainly gabapentin) is employed, but studies evaluating these therapies are uncommon.
- Surgical joint replacement is available for the hip.

References

1. Analysis of normal and osteoarthritic canine cartilage mRNA expression by quantitative polymerase chain reaction. Analysis of normal and osteoarthritic canine cartilage mRNA expression by quantitative polymerase chain reaction. 2011;1–9.
2. Benito J, Hansen B, DePuy V, Davidson GS, Thomson A, Simpson W, Roe S, Hardie E, Lascelles BDX. Feline Musculoskeletal Pain Index: Responsiveness and Testing of Criterion Validity. J Vet Intern Med 2013;27:474–482.
3. Bennett D, Zainal Ariffin SMB, Johnston P. Osteoarthritis in the cat: 2. How should it be managed and treated? Journal of Feline Medicine and Surgery 2012;14:76–84.



© Copyright 2016 International Association for the Study of Pain. All rights reserved.

IASP brings together scientists, clinicians, health-care providers, and policymakers to stimulate and support the study of pain and translate that knowledge into improved pain relief worldwide.

4. Brown DC, Boston RC, Coyne JC. Ability of the canine brief pain inventory to detect response to treatment in dogs with osteoarthritis. *Journal of the American Veterinary Medical Association* 2008.
5. Brown DC, Boston RC, Farrar JT. Use of an activity monitor to detect response to treatment in dogs with osteoarthritis. *Journal of the American Veterinary Medical Association* 2010;237:66–70.
6. Brydges NM, Argyle DJ, Mosley JR, Duncan JC, Fleetwood-Walker S, Clements DN. Clinical assessments of increased sensory sensitivity in dogs with cranial cruciate ligament rupture. *The Veterinary Journal* 2012;193:545–550.
7. Innes JF, Clegg P. Comparative rheumatology: what can be learnt from naturally occurring musculoskeletal disorders in domestic animals? *Rheumatology (Oxford)* 2010;49:1030–1039.
8. Diagnosis and treatment of osteoarthritis. *Diagnosis and treatment of osteoarthritis*. 2010;25:20–25. doi:10.1053/j.tcam.2009.10.005.
9. Freire M, Meuten D, Lascelles D. Pathology of Articular Cartilage and Synovial Membrane From Elbow Joints With and Without Degenerative Joint Disease in Domestic Cats. *Veterinary Pathology* 2014;51:968–978.
10. Guillot M, Taylor PM, Rialland P, Klinck MP, Martel-Pelletier J, Pelletier J-P, Troncy E. Evoked temporal summation in cats to highlight central sensitization related to osteoarthritis-associated chronic pain: a preliminary study. *PLoS ONE* 2014;9:e97347.
11. Initial evaluation of nighttime restlessness in a naturally occurring canine model of osteoarthritis pain. Initial evaluation of nighttime restlessness in a naturally occurring canine model of osteoarthritis pain. 2015;3:e772. doi:10.7717/peerj.772.
12. Lascelles BDX, Freire M, Roe SC, DePuy V, Smith E, Marcellin-Little DJ. Evaluation of Functional Outcome After BFX® Total Hip Replacement Using a Pressure Sensitive Walkway. *Veterinary Surgery* 2010;39:71–77.
13. Lascelles BDX, Gaynor JS, Smith ES, Roe SC, Marcellin-Little DJ, Davidson G, Boland E, Carr J. Amantadine in a Multimodal Analgesic Regimen for Alleviation of Refractory Osteoarthritis Pain in Dogs. *J Vet Intern Med* 2008;22:53–59.
14. Lascelles BDX, Henry JB III, Brown J, Robertson I, Sumrell AT, Simpson W, Wheeler S, Hansen BD, Zamprogno H, Freire M, Pease A. Cross-Sectional Study of the Prevalence of Radiographic Degenerative Joint Disease in Domesticated Cats. *Veterinary Surgery* 2010;39:535–544.
15. Percie du Sert N, Rice ASC. Improving the translation of analgesic drugs to the clinic: animal models of neuropathic pain. *Br J Pharmacol* 2014;171:2951–2963.
16. Vandeweerd JM, Coison C, Clegg P, Cambier C, Pierson A, Hontoir F, Saegerman C, Gustin P, Buczinski S. Systematic Review of Efficacy of Nutraceuticals to Alleviate Clinical Signs of Osteoarthritis. *J Vet Intern Med* 2012;26:448–456.
17. Walton MB, Cowderoy E, Lascelles D, Innes JF. Evaluation of Construct and Criterion Validity for the “Liverpool Osteoarthritis in Dogs” (LOAD) Clinical Metrology Instrument and Comparison to Two Other Instruments. *PLoS ONE* 2013;8:e58125.
18. Williams MD, Kirkpatrick AE, Griffith E, Benito J, Hash J, Lascelles BDX. Feasibility and repeatability of thermal quantitative sensory testing in normal dogs and dogs with hind limb osteoarthritis-associated pain. *Vet. J.* 2014;199:63–67.

About the International Association for the Study of Pain®

IASP is the leading professional forum for science, practice, and education in the field of pain. [Membership is open to all professionals](#) involved in research, diagnosis, or treatment of pain. IASP has more than 7,000 members in 133 countries, 90 national chapters, and 20 Special Interest Groups.

Plan to join your colleagues at the [16th World Congress on Pain](#), September 26-30, 2016, in Yokohama, Japan.

As part of the Global Year Against Pain in the Joints, IASP offers a series of 20 Fact Sheets that cover specific topics related to joint pain. These documents have been translated into multiple languages and are available for free download. Visit www.iasp-pain.org/globalyear for more information.



© Copyright 2016 International Association for the Study of Pain. All rights reserved.

IASP brings together scientists, clinicians, health-care providers, and policymakers to stimulate and support the study of pain and translate that knowledge into improved pain relief worldwide.