



## Prevention of Pain in Vulnerable Populations

The prevention of pain is a global public health concern, and a priority [1] because of the sheer numbers affected. The major cause of pain in all populations is trauma and the persistence of pain following both injury and surgical intervention represents a potentially enormous problem for society.

Vulnerable populations are known to experience significant health disparities [2]. The World Health Organization [3] identifies vulnerable populations as including children, pregnant women, older adults, malnourished people, and people who are ill or immunocompromised. These minority and underserved groups may also be affected by poverty with limited access to healthcare or public health interventions. Consequently, the prevention of pain in vulnerable populations should be viewed as a safeguarding issue.

Additionally, those who are in pain may become vulnerable because of the debilitating effects of pain and suffering. Amongst potentially vulnerable groups, older adults are more likely to experience pain than other segments of the adult population [4]. However, vulnerability is not confined to the oldest in society, and the non-verbal are especially at risk of having their pain poorly assessed or managed [5].

Musculoskeletal conditions including low back and neck pain are major causes of disability in all adult populations [6]. Ageing and disability increases the potential for chronic pain [7]. However, 63% of older adults with dementia had bothersome chronic pain compared to 54% of adults without dementia in a sample of 7,609 community-dwelling older adults [8]. The common pain sites are knees, hips and low back often associated with osteoarthritis and osteoporosis. Painful musculoskeletal problems can increase the risk of cardiovascular disease and for older adults; increase the risk of falls and mortality [7]. Prevention of pain and associated disability can be supported with adopting a healthy lifestyle including regular physical activity [9].

Chronic pain prevention is more challenging. The triggers and pathology of chronic pain are complex; in particular there may be no identifiable causal event [9]. However, there may be particular risk factors associated with the development of chronic pain which could be avoided in particular populations. Psychological factors, such as anxiety, depression, resilience and pain beliefs, are known to be a considerable influence on the experience of chronic pain [10]. Furthermore, older adults have a lower pain threshold which may progressively decrease over time in older adults with dementia, while their pain tolerance increases because they are unable to cognitively recognize and quickly interpret pain, thus increasing vulnerability to consequences of pain [5].

### Recommendations for Prevention of Acute Pain in Vulnerable Populations

- Eat well and maintain a healthy weight to keep bones and muscles working efficiently.
- Keep moving and use exercise to build and maintain core strength and flexibility.

©Copyright 2020 International Association for the Study of Pain. All rights reserved. IASP brings together scientists, clinicians, healthcare providers, and policymakers to stimulate and support the study of pain and translate that knowledge into improved pain relief worldwide.



- Move well; avoid bad posture and over exertion such as heavy lifting wherever possible.
- Reduce stress and anxiety; learn relaxation techniques and coping strategies such as yoga, tai-chi or mindful meditation to stay in control.

### Recommendations for Prevention of Chronic Pain in Vulnerable Populations

All of the recommendations for acute pain prevention plus:

- Effective management of acute pain
- Recognition of chronic pain development as a public health problem

### Assessment and Management

There are times when pain cannot be prevented, and this is when we need to employ effective assessment and management strategies. There are many guidelines that have been developed around the world which can be used to guide the assessment and the management process for all vulnerable populations. At least ten guidelines, for example, focus on older adults and cover issues such as pain assessment and management, acute and chronic pain, or pain associated with osteoarthritis [14, 15, 17, 18, 19, 20, 21].

From a paediatric perspective, guidelines focus on acute, chronic or cancer pain from specific countries and from the World Health Organization [22, 23, 24, 25]. In terms of torture survivors, there are a few guidelines but often focus upon the management of psychological disorders [26] rather than pain. Nevertheless, there have been a number of publications recommendations for managing pain in this population [5].

Organisational and professional barriers often inhibit implementation of these guidelines, so education and improved awareness are the key. Early intervention will prevent development of chronic pain and subsequently reduce the burden upon the patient, their family and society. However, staff need to be aware of how to elicit pain history and responses. For example, in the UK, the 7-minute GP consultation prevents older adults from describing their pain narrative, but it is an organisational limitation that has to be worked around. Similarly, Tai-Seale et al [13] found discussions around pain lasted only 2-3 minutes which is often confounded by communication, gender or cultural factors.

Nevertheless, we are where we are and the organisational, educational barriers are a reality and all we can do is work within the limits of what we have, relying upon the humanity of care to do the best for patients with the tools and guidelines available. At least in 2019/20 we have the skills to recognise the populations who may be considered vulnerable and their specific needs are being highlighted. For example, we have at least 12 pain assessment tools which have been specifically developed for older adults with dementia and some are widely used in clinical practice. For example: PAINAd, PACSLAC, Doloplus, Abbey [14].

## REFERENCES

- [1] Goldberg DS, McGee SJ. Pain as a global public health priority. *BMC public health*. 2011 Dec;11(1):770. <http://www.biomedcentral.com/1471-2458/11/770>
- [2] Campbell LC, Robinson K, Meghani SH, Vallerand A, Schatman M, Sonty N. Challenges and opportunities in pain management disparities research: Implications for clinical practice, advocacy, and policy. *The Journal of Pain*. 2012 Jul 1;13(7):611-9. <https://doi.org/10.1016/j.jpain.2012.02.004>
- [3] World Health Organisation [https://www.who.int/environmental\\_health\\_emergencies/vulnerable\\_groups/en/](https://www.who.int/environmental_health_emergencies/vulnerable_groups/en/)
- [4] Schofield P. *Rev Pain* 2007 Aug; 1(1): 12–14. <https://doi.org/10.1177/204946370700100104>
- [5] International Association for the Study of Pain. Global Year Against pain in vulnerable populations (2019) [https://s3.amazonaws.com/rdcms-iasp/files/production/public/1\\_Guidelines\\_for\\_the\\_Management\\_of\\_Pain\\_in\\_Vulnerable\\_Populationsedited.pdf](https://s3.amazonaws.com/rdcms-iasp/files/production/public/1_Guidelines_for_the_Management_of_Pain_in_Vulnerable_Populationsedited.pdf)
- [6] Booker.SQ., Haedtke (2017) Controlling Pain & Discomfort: Assessment in Non-Verbal older adults. *Nursing*. 2016 May; 46(5): 66–69.
- [7] Molton I, Cook KF, Smith AE, Amtmann D, Chen WH, Jensen MP. Prevalence and impact of pain in adults aging with a physical disability: comparison to a US general population sample. *Clin J Pain*. 2014 Apr; 30(4):307-15. doi: 10.1097/AJP.0b013e31829e9bca.
- [8] Hunt LJ, Covinsky KE, Yaffe K, Stephens CE, Miao Y, Boscardin WJ, Smith AK. (2015) Pain in Community-Dwelling Older Adults with Dementia: Results from the National Health and Aging Trends Study. *J Am Geriatr Soc*. 2015 Aug;63(8):1503-11. doi: 10.1111/jgs.13536. Epub 2015 Jul 22.
- [9] Shega JW, Andrew M, Kotwal A, Lau DT, Herr K, Ersek M, Weiner DK, Chin MH, Dale W. Relationship Between Persistent Pain and 5-Year Mortality: A Population-Based Prospective Cohort Study. *Journal of the American Geriatrics Society*. 2013 Dec;61(12):2135-41.
- [10] Skillgate. E et al (2017) Healthy lifestyle behavior and risk of long duration troublesome neck pain or low back pain among men and women: results from the Stockholm Public Health Cohort. *Clin Epidemiol*. 2017; 9: 491–500.
- [11] Blyth FM, Van Der Windt DA, Croft PR. Chronic disabling pain: a significant public health problem. *American Journal of Preventive Medicine*. 2015 Jul 1;49 (1):98-101.
- [12] Ramírez-Maestre C, Esteve R. The role of sex/gender in the experience of pain: resilience, fear, and acceptance as central variables in the adjustment of men and women with chronic pain. *The Journal of Pain*. 2014 Jun 1;15(6):608-18.
- [13] Tai-Seale.M., Management of chronic pain among older patients: Inside primary care in the US. *European Journal of Pain* 2011. Nov 15(10)
- [14] Schofield PA., et al (2018) The assessment of pain in older people: UK National Guidelines. *Age & Ageing*. Volume 47, Issue suppl\_1, March 2018, Pages i1–i22
- [15] Hadjistavropoulos T. Pain in older persons. *Pain Res Manag*. 2007;12(3):176.
- [16] Rahman MM, Kopec JA, Anis AH, Cibere J, Goldsmith CH. Risk of cardiovascular disease in patients with osteoarthritis: a prospective longitudinal study. *Arthritis Care Res (Hoboken)*. 2013 Dec;65(12):1951-8.
- [17] Singh JA, Furst DE, Bharat A, et al. 2012 update of the 2008 American College of Rheumatology recommendations for the use of disease-modifying antirheumatic drugs and biologic agents in the treatment of rheumatoid arthritis. *Arthritis Care Res (Hoboken)*. 2012;64(5):625–639. doi:10.1002/acr.21641
- [18] American Geriatrics Society 2015 Beers Criteria Update Expert Panel. American Geriatrics Society 2015 Updated Beers Criteria for Potentially Inappropriate Medication Use in Older Adults. *J Am Geriatr Soc*. 2015;63(11):2227–2246.
- [19] Australian and New Zealand College of Anaesthetists. Recommendations on monitoring during anaesthesia. ANZCA 2013. <http://www.anzca.edu.au/documents/ps18-2013-recommendations-on-monitoring-during-ana>
- [20] Arnstein P and Herr KA. Persistent pain management in older adults. The University of Iowa. 2015.
- [21] Cornelius R, Herr KA, Gordon DB, and Kretzer K. Acute pain management in the older adult. The University of Iowa. 2016.
- [22] The Royal College of Emergency Medicine. Best Practice Guideline: Management of Pain in Children. 2017. [https://www.rcem.ac.uk/docs/RCEM%20Guidance/RCEM%20Pain%20in%20Children%20-%20Best%20Practice%20Guidance%20\(REV%20Jul%202017\).pdf](https://www.rcem.ac.uk/docs/RCEM%20Guidance/RCEM%20Pain%20in%20Children%20-%20Best%20Practice%20Guidance%20(REV%20Jul%202017).pdf)
- [23] National Institute of Health & Care Excellence. Chronic pain: assessment and management. 2018. <https://www.nice.org.uk/guidance/gid-ng10069/documents/final-scope>
- [24] World Health Organization. Persisting pain in children package: WHO guidelines on the pharmacological treatment of persisting pain in children with medical illnesses. World Health Organization. 2012. <https://apps.who.int/iris/handle/10665/44540>
- [25] Royal College of Nursing. The recognition and assessment of acute pain in children. 2009. [https://www2.rcn.org.uk/\\_data/assets/pdf\\_file/0004/269185/003542.pdf](https://www2.rcn.org.uk/_data/assets/pdf_file/0004/269185/003542.pdf)
- [26] Amris K and Williams AC. Managing chronic pain in survivors of torture. *Pain Manag*. 2015;5(1):5-12.

©Copyright 2020 International Association for the Study of Pain. All rights reserved. IASP brings together scientists, clinicians, healthcare providers, and policymakers to stimulate and support the study of pain and translate that knowledge into improved pain relief worldwide.



## **AUTHORS**

Margaret Dunham PhD  
Faculty of Health & Wellbeing  
Sheffield Hallam University, UK

Pat Schofield PhD  
Professor of Pain & Ageing  
Sheffield Hallam University, UK

## **REVIEWERS**

Professor Denis Martin  
Centre for Rehabilitation, Exercise and Sports Science  
Teesside University  
Middlesbrough, United Kingdom

Professor Gisèle Pickering, MD, PhD, DPharm,  
Clinical Pharmacology Department  
University Hospital CHU  
Clermont-Ferrand, France

©Copyright 2020 International Association for the Study of Pain. All rights reserved.  
IASP brings together scientists, clinicians, healthcare providers, and policymakers to stimulate and support the study of pain and translate that knowledge into improved pain relief worldwide.

