

Approach to activity, biomechanical loads and flare-ups of back pain

Individual views of patients and their clinicians increasingly play an important role in research. That is especially true for beliefs about the association between physical activity, biomechanical loads and new episodes of back pain. Layperson and clinician views might stimulate research questions and new concepts for improving pain management. Within this context, healthcare providers should be aware of at least 10 facts on pain behavior in patients with back pain:

1. Activity-related pain behavior is highly variable among patients with back pain.

Patients' approaches to physical activities when experiencing pain vary widely. Common approaches include: 1) escape or avoidance of an activity, even at low levels of pain, 2) enduring an activity while tolerating low to moderate levels of pain, 3) persisting with an activity to the point that pain is significantly aggravated (i.e. severe flare-ups of pain), and 4) persisting with an activity until a severe flare-up of pain forces to interrupt or stop this activity, also known as overactivity [7, 18, 26]. Approaches to activity are not always mutually exclusive; individuals can avoid certain activities or movements but persist with other activities to the point of a severe flare-up [3, 4, 27].

2. Patients showing elevated avoidance behavior feel highly disabled by their pain.

The literature provides robust evidence that, in patients with low back pain, avoidance of activities that are thought to increase the experienced level of pain yields a number of detrimental consequences, i.e. that it involves the risk of increasing fear of certain activities or the risk of pain disability in daily life [1].

3. Flare-ups of pain are common among patients with chronic back pain.

Chronic back pain is not of constant and stable intensity but can vary over time with signs of "flare-ups", where pain is much worse than usual for days, weeks, or months [40]. More recently, a multiphase process, including consumers' views and expert consensus extended this definition of a pain flare-up to a worsening of the condition, that is difficult to tolerate and impacts usual activities and/or emotions [13].

4. Physical activities associated with high or sustained biomechanical load are perceived as common triggers of flare-ups.

Patients with chronic back pain [1,12], as well as clinicians [42], perceive physical activities that are associated with high or sustained biomechanical loads (i.e. lifting heavy loads, bending, sustaining longer periods of time in static postures) as the most important precursors of a new acute onset of pain [36, 41] or a flare-up of chronic pain. Objectively assessed static postures (upright or forward bent sitting or standing) in daily life are positively related with pain intensity and pain-related fatigue in patients with chronic back pain [37]. Moreover, results from systematic reviews and meta-analyses give support for the association of biomechanical stressors such as lifting [11], bending [45], and new episodes of back pain, with some studies shown a dose-response [30].

5. Research suggests a time delay between engaging in physical activity and a flare-up of pain.

An individual's perception of an association of biomechanical triggers and a flare-up of pain can be hampered by a delay that exists between biomechanical loadings and an exacerbation of pain. Preliminary studies have shown that dysfunctional physical activities can be followed by a flare-up of pain with a delay of 30 minutes up to hours [20, 32, 36].

6. Pain persistence and overactivity are common behavioral responses to LBP.

Besides avoidance, persistence in an activity despite severe levels of pain is a common behavioral response to musculoskeletal and back pain [1, 9, 14, 17, 19, 22, 25, 27, 29, 31, 33, 34, 39]. Most studies assessed self-reported frequency of persistence behavior, using questionnaires, such as the Avoidance-Endurance Questionnaire AEQ [25], the 9-item Avoidance-Endurance Fast Screen (AEFS) [47], or the Patterns of Activity Measure-Pain POAM-P [8]. Different terms such as pain persistence [17, 25, 27, 39], overdoing [9, 34], task persistence [31] or pain-related endurance [25, 27] have been used to denote persisting with activities in spite of severe pain. In contrast, the terms overactivity [3, 7] or excessive persistence [31] refer to a process of pain persistence that was only halted by an intolerable intensity of pain and a subsequent phase of functional incapacity [28]. The Overactivity in Persistent Pain Assessment (OPPA) is a self-report measure to assess not only persistence despite pain but also the subsequent phase of physical inactivity [46].

7. While adaptivity of pure frequency of pain persistence is inconclusive, habitual overactivity has been associated with worse outcomes.

The term pain persistence behavior that merely refers to the frequency or severity of occurrence displays low or zero linear correlations with pain intensity and inconsistent results with positive or negative relations to pain disability.^{1,14,25,33} Negative associations with low to moderate effect sizes were shown for pain persistence and psychological distress, such as depression, anxiety, pain catastrophizing or fear of movement.^{1,25,33,39} In contrast, measures that include aspects of the process of overactivity, which is followed by an interruption of an activity due to intolerable pain revealed positive associations with disability or psychological distress.^{7,10,31,46}

8. Research has revealed valuable insights into complex patterns of cognitive-affective and behavioral responses to pain.

Inspecting more complex individual patterns of cognitive, affective and behavioral pain responses instead of the frequency of pain persistence, provide more insight into physical and psychological adaptation. For example, patients showing a pattern of distress-endurance pain responses (DER) with negative mood, elevated thought suppression and pain persistence behavior reported significantly higher pain intensity, disability and poorer psychological functioning than those patients with an adaptive response pattern (AR, i.e. with low persistence, low thought suppression and low negative mood) [22, 27, 43]. Despite higher pain and disability, DER patients have been shown to display a significantly higher number of static postures, assessed objectively with a tri-axial accelerometer advice, compared to AR patients [24, 37]. Conversely, patients with an eustress-endurance pattern (EER) with high pain persistence behavior, positive mood despite pain and elevated cognitive distraction from pain displayed higher pain intensity scores than AR patients, but comparably low disability and positive psychological function. Pain cognitions and affective responses seem to determine whether pain persistence is positive or negatively related

to disability and poor psychological functioning. The existence of two different pain persistence groups (similar to DER and EER) besides patients with a fear-avoidance (FAR, high fear of pain or injury, high avoidance behavior) or an adaptive pattern are described in a number of studies [8, 17, 29, 34, 37, 38, 47, 48].

9. Neurobehavioral consequences of different approaches to physical activity requires further research.

The Avoidance-Endurance model (AEM) of pain conceptualizes possible neurobehavioral mechanisms suggesting that, due to extreme pain persistence behavior, these patients are prone to physical overuse/overload and the induction of early muscle fatigue, reduced blood/ oxygen supply, small and repetitive damages of soft tissues, such as muscles, ligaments, and tendons [26, 28]. Conversely, patients with an elevated FAR pattern run the risk of physical de-conditioning, including negative changes in muscular, motor, cardio-respiratory and metabolic aspects of physical fitness [44]. Research that addresses adaptations in the motor control system due to pain persistence and overactivity is currently at the stage of hypotheses, however a number of physiological mechanisms have been outlined suggesting promising perspectives for future studies [28].

10. Flexible and self-determined pacing behavior might be an adaptive response to pain but more research is needed.

Activity pacing refers to regulating activity levels and/or rate in the service of an adaptive goal [35]. Activity pacing is a key self-management strategy taught in pain management programs across the globe where individuals are taught to break up and reschedule pain provoking activities by taking short rest breaks or alternating activities/postures [6]. Outcome studies evaluating the effectiveness of pacing as an intervention are however sparse with inconsistent findings observed.²¹ Qualitative data has highlighted the potential value of activity pacing as a treatment strategy⁴ and activity patterns consistent with the effective use of activity pacing strategies have been associated with better outcomes in some studies. For example, patients with an adaptive pain response pattern (suggestive of an effective pacing pattern [26]) showed the lowest pain intensity and disability scores [27, 29, 43] and lower depression than patients with an avoidance or persistence pattern [9, 27, 29]. In contrast, research on the linear association between frequency of pacing behavior and pain, disability or psychological function have yielded highly inconsistent results [1, 9, 10, 14, 15, 16, 31]. Further research on the effectiveness of activity pacing as a treatment strategy and exploring activity patterns consistent with the effective use of pacing strategies is warranted.

REFERENCES

- [1] Andrews NE, Strong J, Meredith PJ. Activity pacing, avoidance, endurance, and associations with patient functioning in chronic pain: A systematic review and meta-analysis. *Arch Phys Med Rehabil* 2012; 93: 2109–2121.
- [2] Andrews NE, Strong J, Meredith PJ, D'Arrigo RG. Association between physical activity and sleep in adults with chronic pain: A momentary, within-person perspective. *Phys Ther* 2014; 94:499–510.
- [3] Andrews NE, Strong J, Meredith PJ. Overactivity in chronic pain: Is it a valid construct? *Pain* 2015; 156: 1991–2000.
- [4] Andrews NE, Strong J, Meredith PJ, Gordon K, Bagraith, K. "It's very hard to change yourself": an exploration of overactivity in people with chronic pain using an interpretative phenomenological analysis. *Pain* 2015; 156 (7): 1215-1231.
- [5] Andrews NE, Strong J, Meredith PJ, Fleming, JA. The relationship between overactivity and opioid use in chronic pain: A 5-day observational study. *Pain* 2016; 157: 466–474.
- [6] Andrews NE, Deen M. Defining activity pacing: is it time to jump off the merry-go-round?. *J Pain* 2016; 17 (12); 1359-1362.
- [7] Andrews NE, Chien C-W, Ireland D, Varnfield M. Overactivity assessment in chronic pain: the development and psychometric evaluation of a multifaceted self-report assessment. *Eur J Pain* 2020; epub ahead of print. <https://doi.org/10.1002/ejp.1664>
- [8] Cane DB, Nielson WR, McCarthy M, Mazmanian D. Pain-related activity patterns: measurement, interrelationships, and associations with psychosocial functioning. *Clin J Pain* 2013;29:435–442.
- [9] Cane D, Nielson WR, Mazmanian D. Patterns of pain-related activity: replicability, treatment-related changes, and relationship to functioning. *Pain* 2018;159:2522–2529.
- [10] Cane D, Mazmanian D. Development and Initial evaluation of the Patterns of Activity Measure-Pain Short Form. *Clin J Pain* 2020;36:675-682.
- [11] Coenen P, Gouttebarga V, Van der Burght ASAM, Van Dieen JH, Frings-Dresen MHW, Van der Beek AJ, Burdorf A. The effect of lifting during work on low back pain: a health impact assessment based on a meta-analysis. *Occup Environ Med* 2014;71:871–877
- [12] Costa N, Hodges PW, Ferreira ML, Makovey J, Setchell J. What Triggers an LBP Flare? A Content Analysis of Individuals' Perspectives. *Pain Med* 2020;21:13-20
- [13] Costa N, Ferreira ML, Setchell J, Makovey J, Dekroot T, MChir AD, Diwan A, Koes B, Natvig B, Vocenzino B, Hunter D, Roseen E, Rasmussen-Barr, Guillemin F, Hartvigsen J, Bennell K, Costa L, Macedo L, Pinheiro M, Underwood M, Van Tulder M, Johansson M, Enthoven P, Kent P, O'Sullivan P, Suri P, Genevay S, Hodges PW. A definition of flare in low back pain (LBP): A multiphase process involving perspectives of individuals with LBP and expert consensus. *J Pain acc paper* <https://doi.org/10.1016/j.jpain.2019.03.009>
- [14] Esteve R, Ramirez-Maestre ML, Peters ER, SerranoIbanez GT, Ruiz-Parraga GT, Lopez-Martinez AE. Development and initial validation of the activity patterns scale in patients with chronic pain. *J Pain* 2016;17:451– 461, 2016.
- [15] Esteve R, Lopez-Martinez AE, Peters ML, Serrano-Ibanez ER, RuizParraga GT, Gonzalez-Gomez H, Ramirez- Maestre C. Activity pattern profiles: relationship with affect, daily functioning, impairment, and variables related to life goals. *J Pain* 2017;18:546–55.
- [16] Esteve R, Lopez-Martinez AE, Peters ML, Serrano-Ibanez ER, Ruiz-Parraga GT, Ramirez-Maestre C. Optimism, positive and negative affect, and goals adjustment strategies: their relationship to activity patterns in patients with chronic musculoskeletal pain. *Pain Res Manage* 2018;Article ID 6291719 <https://doi.org/10.1155/2018/6291719>
- [17] Fehrmann E, Tuechler K, Kienbacher T, et al. Comparisons in muscle function and training rehabilitation outcomes between avoidance-endurance model-subgroups. *Clin J Pain* 2017;33:912–920.
- [18] Fordyce WE. *Behavioral Methods for Chronic Pain and Illness*. St. Louis, MO: Mosby, 1976.
- [19] Gajsar H, Titze C, Levenig C, Kellmann M, Heidari J, Kleinert J, Rusu AC, Hasenbring MI. Psychological pain responses in athletes and non-athletes with low back pain Avoidance and endurance matter. *Eur J Pain* 2019;23: 1649-1662.
- [20] Geisser ME, Robinson ME, Richardson C. A time series analysis of the relationship between ambulatory EMG, pain, and stress in chronic low back pain. *Biofeedback Self Regul* 1995;20:339–355.
- [21] Guy, L., C. McKinstry, and C. Bruce, Effectiveness of Pacing as a Learned Strategy for People With Chronic Pain: A Systematic Review. *The American Journal Of Occupational Therapy*, 2019. 73(3): p. 7303205060p1- 7303205060p10.
- [22] Hasenbring M. Endurance strategies-a neglected phenomenon in the research and therapy of chronic pain? *Schmerz* 1993;7:304–313.
- [23] Hasenbring M, Ulrich HW, Hartmann M, Soyka D. The efficacy of a risk factorbased cognitive behavioral intervention and electromyographic biofeedback in patients with acute sciatic pain. An attempt to prevent chronicity. *Spine (Phila Pa 1976)* 1999;24:2525–35.
- [24] Hasenbring MI, Plaas H, Fischbein B, Willburger R. The relationship between activity and pain in patients 6 months after lumbar disc surgery: do painrelated coping modes act as moderator variables? *Eur J Pain* 2006;10:701–9.
- [25] Hasenbring MI, Hallner D, Rusu AC. Fear-avoidance- and endurance-related responses to pain: development and validation of the Avoidance-Endurance Questionnaire (AEQ). *Eur J Pain* 2009;13:620–8.

- [26] Hasenbring MI, Verbunt JA. Fear-avoidance and endurance-related responses to pain: new models of behavior and their consequences for clinical practice. *Clin J Pain* 2010;26:747–53.
- [27] Hasenbring MI, Hallner D, Klasen B, et al. Pain-related avoidance versus endurance in primary care patients with subacute back pain: psychological characteristics and outcome at a 6-month follow-up. *Pain* 2012;153:211–217.
- [28] Hasenbring MI, Andrews NE, Ebenbichler G. Overactivity in chronic pain, the role of pain-related endurance and neuromuscular activity. *Clin J Pain* 2020;36:162–171.
- [29] Huijinen IPJ, Verbunt JA, Peters ML, Smeets RJEM, Kindermans HPJ, Roelofs J, Goossens M, Seelen HAM. Differences in activity-related behaviour among patients with chronic low back pain. *Eur J Pain* 2011;15:748–55.
- [30] Jansen JP, Morgenstern H, Burdorf A. Dose-response relations between occupational exposures to physical and psychosocial factors and the risk of low back pain. *Occup Environ Med* 2004;61:972–979.
- [31] Kindermans HPJ, Roelofs J, Goossens MEJB, et al. Activity patterns in chronic pain: underlying dimensions and associations with disability and depressed mood. *J Pain*. 2011;12:1049–1058.
- [32] Liszka-Hackzell JJ, Martin DP. An analysis of the relationship between activity and pain in chronic and acute low back pain. *Anesth Analg* 2004;99:477–481.
- [33] Luthi F, Vuistiner P, Favre C, et al. Avoidance, pacing, or persistence in multidisciplinary functional rehabilitation for chronic musculoskeletal pain: an observational study with cross-sectional and longitudinal analyses. *PLoS One*. 2018;13: e0203329.
- [34] McCracken LM, Samuel VM. The role of avoidance, pacing, and other activity patterns in chronic pain. *Pain* 2007;130:119–125.
- [35] Nielson WR, Jensen MP, Karsdorp PA, Vlaeyen JW. (2013). Activity pacing in chronic pain: concepts, evidence, and future directions. *Clin J Pain*, 29, pp. 461–468.
- [36] Parreira Pdo C, Maher CG, Latimer J, et al. Can patients identify what triggers their back pain? Secondary analysis of a case-crossover study. *Pain* 2015;156(10):1913–1919.
- [37] Plaas H, Sudhaus S, Willburger R, et al. Physical activity and low back pain: the role of subgroups based on the avoidance-endurance model. *Disabil Rehabil* 2014;36:749–755.
- [38] Scholich SL, Hallner D, Wittenberg RH, et al. Pilot study on pain response patterns in chronic low back pain. The influence of pain response patterns on quality of life, pain intensity and disability. *Schmerz*. 2011;25:184–190.
- [39] Scholich, S. L., Hallner, D., Wittenberg, R. H., Hasenbring, M. I., & Rusu, A. C. (2012). The relationship between pain, disability, quality of life and cognitive-behavioural factors in chronic back pain. *Disability and Rehabilitation* 2012; 34:1993–2000.
- [40] Suri P, Saunders KW, Von Korff M. Prevalence and characteristics of flare-ups of chronic nonspecific back pain in primary care: A telephone survey. *Clin J Pain* 2012;28(7):573–80.
- [41] Steffens D, Ferreira ML, Latimer J, et al. What triggers an episode of acute low back pain? A case-crossover study. *Arthritis Care Res* 2015;67 (3):403–10.
- [42] Stevens ML, Steffens D, Ferreira ML, Latimer J, Blyth F, Maher CG. Patients' and physiotherapists' views on triggers for low back pain. *Spine* 2016;41:E218–E224
- [43] Titz C, Fett D, Trompeter K, Platen P, Gajsar H, Hasenbring MI. Psychosocial subgroups in high-performance-athletes with low back pain: eustress-endurance is most frequent, distress-endurance most problematic! *Scand J Pain* 2020 <https://doi.org/10.1515/sjpain-2020-0053>
- [44] Verbunt JA, Seelen HA, Vlaeyen JW, et al. Disuse and deconditioning in chronic low back pain: concepts and hypotheses on contributing mechanisms. *Eur J Pain* 2003;7:9–21.
- [45] Wai, E.K., Roffey, D.M., Bishop, P., Kwon, B.K., Dagenais, S., 2010. Causal assessment of occupational bending or twisting and low back pain: results of a systematic review. *Spine J*. 10, 76–88.
- [46] Andrews NE, Chien CW, Ireland D, Varnfield M. Overactivity assessment in chronic pain: The development and psychometric evaluation of a multifaceted self-report assessment. *Eur J Pain* 2021;25:225–242.
- [47] Wolff, SV, Willburger R., Hallner D, Rusu AC, Rusche H, Schulte T, Hasenbring MI. Avoidance-endurance fast screening (AEFS). Content and predictive validity of a 9-item screening instrument for patients with unspecific subacute low back pain. *Schmerz* 2020; 34:S1–S7.
- [48] Titz C, Hasenbring MI, Kristensen L, Bendix L, Vaegter HB. Patterns of approach to activity in 851 patients with severe chronic pain: translation and preliminary validation of the 9-item Avoidance-Endurance Fast-Screen (AEFS) into Danish. *Clin J Pain* 2021 DOI:10.1097/AJP.0000000000000912

AUTHORS

Prof. Dr. Monika I Hasenbring
Department of Medical Psychology and Medical Sociology, Faculty of Medicine, Ruhr-University of Bochum, Germany
Faculty of Health Science, University of Southern Denmark, Odense, Denmark
Monika.Hasenbring@ruhr-uni-bochum.de

Nicole E. Andrews
The Occupational Therapy Department and the Professor Tess Cramond Multidisciplinary Pain Centre, The Royal Brisbane and Women's Hospital, Australia
RECOVER Injury Research Centre, The University of Queensland, Australia

Gerold Ebenbichler
Department of Physical Medicine, Rehabilitation and Occupational Medicine,
Vienna Medical University, General Hospital of Vienna, Austria

REVIEWERS

Prof. Dr. Jaap H. van Dieën
Professor of Biomechanics and Head, Department of Human Movement Sciences
VU Amsterdam, Netherlands

Pradeep Suri, MD
Associate Professor and Physician, Department of Rehabilitation Medicine
University of Washington, USA